

Microbiological Safety of Fresh and Fresh-Cut Melons: A Bibliography

Compiled by

Robert L. Buchanan, Ph.D.
Center for Food Safety and Security Systems
College of Agriculture and Natural Resources
University of Maryland
0119 Symons Hall
College Park, Maryland, USA

September, 2009

Golden, D.A. et al. Growth of *Salmonella* spp. in cantaloupe, watermelon, and honeydew melons. *Journal of Food Protection* 56:194-196. 1993.

Castillo, A. and Escartin, E.F. Survival of *Campylobacter jejuni* on sliced watermelon and papaya. *Journal of Food Protection* 57:166-168. 1994.

Herwaldt, B.L. et al. Characterization of a variant strain of Norwalk virus from a food-borne outbreak of gastroenteritis on a cruise ship in Hawaii. *Journal of Clinical Microbiology* 32:861-866. 1994.

Del Rosariao, B.A. and Beuchat, L.R. Survival and growth of enterohemorrhagic *Escherichia coli* O157:H7 in cantaloupe and watermelon. *Journal of Food Protection* 58:105-107. 1995.

Beuchat, L.R. Pathogenic microorganisms associated with fresh produce. *Journal of Food Protection* 59:204-216. 1996.

Brackett, R.E. Incidence, contributing factors, and control of bacterial pathogens in produce. *Postharvest Biology and Technology* 15:305-311. 1999.

Larson, A.E. and Johnson, E.A. Evaluation of botulinal toxin production in packaged fresh-cut cantaloupe and honeydew melons. *Journal of Food Protection* 62:948-952. 1999.

Park, C.M. and Beuchat, L.R. Evaluation of sanitizers for killing *Escherichia coli* O157:H7, *Salmonella*, and naturally occurring microorganisms on cantaloupes,

honeydew melons, and asparagus. *Dairy, Food and Environmental Sanitation* 19:842-847. 1999.

Leverentz, B. et al. Examination of bacteriophage as a biocontrol method for *Salmonella* on fresh-cut fruit: A model study. *Journal of Food Protection* 64:1116-1121. 2001.

Sapers, G.M. et al. Antimicrobial treatments for minimally processed cantaloupe melon. *Journal of Food Science* 66:345-349. 2001.

Shearer, A.E.H. et al. Evaluation of a polymerase chain reaction-based system for detection of *Salmonella* Enteritidis, *Escherichia coli* O157:H7, *Listeria* spp., and *Listeria monocytogenes* on fresh fruits and vegetables. *Journal of Food Protection* 64:788-795. 2001.

Ukuku, D.O. and Sapers, G.M. Effect of sanitizer treatments on *Salmonella* Stanley attached to the surface of cantaloupe and cell transfer to fresh-cut tissues during cutting practices. *Journal of Food Protection* 64:1286-1291. 2001.

Ukuku, D.O. et al. Influence of washing treatment on native microflora and *Escherichia coli* population of inoculated cantaloupes. *Journal of Food Safety* 21:31-47. 2001.

Ukuku, D.O. et al. Bioluminescence ATP assay for estimating total plate counts of surface microflora of whole cantaloupe and determining efficacy of washing treatments. *Journal of Food Protection* 64:813-819. 2001.

Anderson, S.M. et al. Multistate outbreaks of *Salmonella* serotype Poona infections associated with eating cantaloupe from Mexico --- United States and Canada, 2000—2002. *Morbidity and Mortality Weekly Report* 51:1044-1047. 2002.

Kozempel, M. et al. Application of the vacuum/steam/vacuum surface intervention process to reduce bacteria on the surface of fruits and vegetables. *Innovative Food Science and Emerging Technologies* 3:62-72. 2002.

Roller, S. and Seedhar, P. Carvacol and cinnamic acid inhibit microbial growth in fresh-cut melon and kiwifruit at 4° and 8°C. *Letters in Applied Microbiology* 35:390-394. 2002.

Ukuku, D.O. and Fett, W.F. Effectiveness of chlorine and nisin-EDTA treatments of whole melons and fresh-cut pieces for reducing native microflora and extending shelf-life. *Journal of Food Safety* 22:231-253. 2002.

Ukuku, D.O. and Fett, W.F. Behavior of *Listeria monocytogenes* inoculated on cantaloupe surfaces and efficacy of washing treatments to reduce transfer from rind to fresh-cut pieces. *Journal of Food Protection* 65:924-930. 2002.

- Barak, J.D. et al. Recovery of surface bacteria from and surface sanitization of cantaloupes. *Journal of Food Protection* 66:1805-1810. 2003.
- Caldwell, K.N. et al. Attraction of a free-living nematode, *Caenorhabditis elegans*, to foodborne pathogenic bacteria and its potential as a vector of *Salmonella* Poona for preharvest contamination of cantaloupe. *Journal of Food Protection* 66:1964-1971. 2003.
- Gagliardi, J.V. et al. On-farm and postharvest processing of bacterial contamination to melon rinds. *Journal of Food Protection* 66:82-87. 2003.
- Leverentz, B. et al. Biocontrol of *Listeria monocytogenes* on fresh-cut produce by treatment with lytic bacteriophages and a bacteriocin. *Applied and Environmental Microbiology* 69:4519-4526. 2003.
- Materon, L.A. Survival of *Escherichia coli* O157:H7 applied to cantaloupes and the effectiveness of chlorinated water and lactic acid as disinfectants. *World Journal of Microbiology and Biotechnology* 19:867-873. 2003.
- Suslow, T.V. et al. Production practices as risk factors in microbial food safety of fresh and fresh-cut produce. *Comprehensive Reviews in Food Science and Food Safety* 2(supplement):38-77. 2003.
- Annous, B.A. et al. Surface pasteurization of whole fresh cantaloupes inoculated with *Salmonella* Poona or *Escherichia coli*. *Journal of Food Protection* 67:1876-1885. 2004.
- Castillo, A. et al. *Salmonella* contamination during production of cantaloupe: A binational study. *Journal of Food Protection* 67:713-720. 2004.
- Hammack, T.S. et al. Relative effectiveness of the Bacteriological Analytical Manual method for the recovery of *Salmonella* from whole cantaloupes and cantaloupe rinses with selected preenrichment media and rapid methods. *Journal of Food Protection* 67:870-877. 2004.
- Karenlampi, R. and Hanninen, M.-L. Survival of *Campylobacter jejuni* on various fresh produce. *International Journal of Food Microbiology* 97:187-195. 2004.
- Leverentz, B. et al. Optimizing concentration and timing of a phage spray application to reduce *Listeria monocytogenes* on honeydew melon tissue. *Journal of Food Protection* 67:1682-1686. 2004.
- Li, Y. and Mustapha, A. Simultaneous detection of *Escherichia coli* O157:H7, *Salmonella*, and *Shigella* in apple cider and produce by a multiplex PCR. *Journal of Food Protection* 67:27-33. 2004.

- Penteado, A.L. and Leitao, M.F.F. Growth of *Listeria monocytogenes* in melon, watermelon, and papaya pulps. *International Journal of Food Microbiology* 91:89-94. 2004.
- Richards, G.M. and Beuchat, L.R. Attachment of *Salmonella* Poona to cantaloupe rind and stem scar tissues as affected by temperature of fruit and inoculum. *Journal of Food Protection* 67:1359-1364. 2004.
- Richards, G.M. et al. Survey of yeasts for antagonistic activity against *Salmonella* Poona in cantaloupe juice and wounds in rinds coinfecting with phytopathogenic molds. *Journal of Food Protection* 67:2132-2142. 2004.
- Rodgers, S.L. et al. A comparison of different chemical sanitizers for inactivating *Escherichia coli* O157:H7 and *Listeria monocytogenes* in solution and on apples, lettuce, strawberries, and cantaloupe. *Journal of Food Protection* 67:721-731. 2004.
- Sanglay, G.C. et al. Recovery of *Salmonella* spp. From raw produce surfaces using ultrasonication. *Foodborne Pathogens and Disease* 1:295-299. 2004.
- Taitt, C.R. et al. Detection of *Salmonella enterica* serovar Typhimurium by using a rapid, array-based immunosensor. *Applied and Environmental Microbiology* 70:152-158. 2004.
- Ukuku, D.O. Effect of hydrogen peroxide treatment on microbial quality and appearance of whole and fresh-cut melons contaminated with *Salmonella* spp. *International Journal of Food Microbiology* 95:137-146. 2004.
- Ukuku, D.O. and Fett, W.F. Effect of nisin in combination with EDTA, sodium lactate, and potassium sorbate for reducing *Salmonella* on whole and fresh-cut cantaloupe. *Journal of Food Protection* 67:2143-2150. 2004.
- Ukuku, D.O. et al. Method of applying sanitizers and sample preparation affects recovery of native microflora and *Salmonella* on whole cantaloupe surfaces. *Journal of Food Protection* 67:999-1004. 2004.
- Ukuku, D.O. et al. Effect of hot water and hydrogen peroxide treatments on survival of *Salmonella* and microbial quality of whole and fresh-cut cantaloupe. *Journal of Food Protection* 67:432-437. 2004.
- Annous, B.A. et al. Improved recovery procedure for evaluation of sanitizer efficacy in disinfecting contaminated cantaloupes. *Journal of Food Science* 70:M242-M247. 2005.
- Annous, B.A. et al. Biofilm formation by *Salmonella* spp. on cantaloupe melon. *Journal of Food Safety* 25:276-287. 2005.

Duffy, E.A. et al. Concentrations of *Escherichia coli* and genetic diversity and antibiotic resistance profiling of *Salmonella* isolated from irrigation water, packaging shed equipment, and fresh produce in Texas. *Journal of Food Protection* 68:70-79. 2005.

Johnston, L.M. et al. A field study of the microbiological quality of fresh produce. *Journal of Food Protection* 69:1840-1847. 2005.

Kim, H. and Beuchat, L.R. Survival and growth of *Enterobacter sakazakii* on fresh-cut fruits and vegetables and in unpasteurized juices as affected by storage temperature. *Journal of Food Protection* 68:2541-2552. 2005.

Parnell, T.L. et al. Reducing *Salmonella* on cantaloupes and honeydew melons using practices applicable to postharvest handling, foodservice, and consumer preparation. *International Journal of Food Microbiology* 99:59-70. 2005.

Richards, G.M. and Beuchat, L.R. Infection of cantaloupe rind with *Cladosporium caldosporioides* and *Penicillium expansum*, and associated migration of *Salmonella* Poona in edible tissue. *International Journal of Food Microbiology* 103:1-10. 2005.

Sharma, M. et al. Thermal tolerance of acid adapted and unadapted *Salmonella*, *Escherichia coli* O157:H7, and *Listeria monocytogenes* in cantaloupe juice and watermelon juice. *Letters in Applied Microbiology* 41:448-453. 2005.

Stine, S.W. et al. Effect of relative humidity on preharvest survival of bacterial and viral pathogens on the surface of cantaloupe, lettuce, and bell peppers. *Journal of Food Protection* 68:1352-1358. 2005.

Stine, S.W. et al. Application of microbial risk assessment to the development of standards for enteric pathogens in water used to irrigate fresh produce. *Journal of Food Protection* 68:913-918. 2005.

Ukuku, D.O. et al. Use of hydrogen peroxide in combination with nisin, sodium lactate and citric acid for reducing transfer of bacterial pathogens from whole melon surfaces to fresh-cut pieces. *International Journal of Food Microbiology* 104:225-233. 2005.

Ukuku, D.O. et al. ATP bioluminescence assay for estimation of microbial populations of fresh-cut melon. *Journal of Food Protection* 68:2427-2432. 2005.

Vargas, A.M. et al. Detection of fecal contamination on cantaloupes using hyperspectral fluorescence imagery. *Journal of Food Science* 70:E471-E476. 2005.

Boynton, B.B. et al. Effects of low-dose electron beam irradiation on respiration, microbiology, texture, color, and sensory characteristics of fresh-cut cantaloupe stored in modified atmosphere packages. *Journal of Food Science* 71:S149-S155. 2006.

- Dipersio, P.A. et al. Sensory evaluation of home dried fruit prepared using treatments that enhance destruction of pathogenic bacteria. *Journal of Food Quality* 29:47-64. 2006.
- Espinoza-Medina, I.E. et al. PCR identification of *Salmonella*: Potential contamination sources from production and postharvest handling of cantaloupes. *Journal of Food Protection* 69:1422-1425. 2006.
- Fan, X. et al. Combination of hot-water surface pasteurization of whole fruit and low-dose gamma irradiation of fresh-cut cantaloupe. *Journal of Food Protection* 69:912-919. 2006.
- Hammack, T.S. et al. Effect of sample preparation and preenrichment media on the recovery of *Salmonella* from cantaloupes, mangoes, and tomatoes. *Journal of AOAC International* 89:180-184. 2006.
- Solomon, E.B. et al. Thermal inactivation of *Salmonella* on cantaloupes using hot water. *Journal of Food Science* 71:M25-M30. 2006.
- Ukuku, D.O. and Fett, W.F. Effects of cell surface charge and hydrophobicity on attachment of 16 *Salmonella* serovars to cantaloupe rind and decontamination with sanitizers. *Journal of Food Protection* 69:1835-1843. 2006.
- Ukuku, D.O. et al. Effect of vacuum-steam-vacuum treatment of microbial quality of whole and fresh-cut cantaloupe. *Journal of Food Protection* 69:1623-1629. 2006.
- Wang, H. et al. Dual-phasic inactivation of *Escherichia coli* O157:H7 with peroxyacetic acid, acidic electrolyzed water and chlorine on cantaloupes and fresh-cut apples. *Journal of Food Safety* 26:335-347. 2006.
- Wang, Z. et al. Influence of gamma irradiation of enzyme, microorganism, and flavor of cantaloupe (*Cucumis melo* L.) juice. *Journal of Food Science* 71:M215-M220. 2006.
- Zhou, B. et al. 1-Methylcyclopropene counteracts ethylene-induced microbial growth on fresh-cut watermelon. *Journal of Food Science* 71:M180-M184. 2006.
- Critzer, F.J. et al. Atmospheric plasma inactivation of foodborne pathogens on fresh produce surfaces. *Journal of Food Protection* 70 :2290-2296. 2007.
- Materon, L.A. et al. Identification of sources of microbial pathogens on cantaloupe rinds from pre-harvest to post-harvest operations. *World Journal of Microbiology and Biotechnology* 23:1281-1287. 2007.
- Mosqueda-Melgar, J. et al. Influence of treatment time and pulse frequency on *Salmonella* ENTERITIDIS, *Escherichia coli* and *Listeria monocytogenes* populations

inoculated in melon and watermelon juices treated by pulsed electric. *International Journal of Food Microbiology* 117:192-200. 2007.

Ukuku, D.O. and Sapers, G.M. Effect of time before storage and storage temperature on survival of *Salmonella* inoculated on fresh-cut melons. *Food Microbiology* 24:288-295. 2007.

Ailes, E.C. et al. Microbial concentrations on fresh produce are affected by postharvest processing, importation, and season. *Journal of Food Protection* 71:2389-2397. 2008.

Akins, E.D. et al. Washing practices on the microflora on Georgia-grown cantaloupes. *Journal of Food Protection* 71:46-51. 2008.

Bassett, J. and McClure, P. A risk assessment approach for fresh fruits. *Journal of Applied Microbiology* 104:925-943. 2008.

Belletti, N. et al. Antimicrobial efficacy of citron essential oil on spoilage and pathogenic microorganisms in fruit-based salads. *Journal of Food Science* 73:M331-M338. 2008.

Fan, X. et al. Effect of hot water surface pasteurization of whole fruit on shelf-life and quality of fresh-cut cantaloupe. *Journal of Food Science* 73:M91-M98. 2008.

Gallegos-Robles, M.A. et al. Identification of *Salmonella* serotypes isolated from cantaloupe and chile pepper production systems in Mexico by PCR–restriction fragment length polymorphism. *Journal of Food Protection* 71:2217–2222. 2008.

Mosqueda-Melgar, J. et al. Combination of high-intensity pulsed electric fields with natural antimicrobials to inactivate pathogenic microorganisms and extend the shelf-life of melon and watermelon juices. *Food Microbiology* 25:479-491. 2008.

Mahmoud, B.S.M. et al. Inactivation kinetics of inoculated *Escherichia coli* O157:H7, *Listeria monocytogenes*, and *Salmonella* Poona on whole cantaloupes by chlorine dioxide gas. *Food Microbiology* 25:857-865. 2008.

Oms-Oliu, G. et al. Effect of superatmospheric and low oxygen modified atmospheres on shelf-life extension of fresh-cut melon. *Food Control* 19:191-199. 2008.

Perni, S. et al. Cold atmosphere plasma decontamination of the pericaps of fruit. *Journal of Food Protection* 71:302-308. 2008.

Perni, S. et al. Cold atmospheric plasma disinfection of cut fruit surfaces contaminated with migrating microorganisms. *Journal of Food Protection* 71:1619-1625. 2008.

Raybaudi-Massilia, R.M. et al. Edible alginate-based coating as carrier of antimicrobials to improve shelf-life and safety of fresh-cut melon. *International Journal of Food Microbiology* 121:313-327. 2008.

Selma, M.V. et al. Effect of gaseous ozone and hot water on microbial and sensory quality of cantaloupe and potential transference of *Escherichia coli* O157:H7 during cutting. *Food Microbiology* 25:162-168. 2008.

Selma, M.V. et al. Reduction by gaseous ozone of *Salmonella* and microbial flora associated with fresh-cut cantaloupe. *Food Microbiology* 25:558-565. 2008.

Silveira, A.C. et al. Alternative sanitizers to chlorine for use on fresh-cut “Galia” (*Cucumis melo* var. *catalupensis*) melon. *Journal of Food Science* 73:M405-M411. 2008.

Hanning, I.B. et al. Salmonellosis outbreaks in the United States due to fresh produce: Sources and potential intervention measures. *Foodborne Pathogens and Disease* 6:635-648. 2009.

Munnoch, S. A. et al. A multi-state outbreak of *Salmonella* Saintpaul in Australia associated with cantaloupe consumption. *Epidemiology and Infection* 137:367-374. 2009.

Sharma, M. et al. Effectiveness of bacteriophages in reducing *Escherichia coli* O157:H7 on fresh-cut cantaloupes and lettuce. *Journal of Food Protection* 72:1481-1485. 2009.

Silagyi, K. et al. Production of biofilm and quorum sensing by *Escherichia coli* O157:H7 and its transfer from contact surfaces to meat, poultry, ready-to-eat deli, and produce products. *Food Microbiology* 26:514-519. 2009

Wang, H. et al. Effect of surface roughness on retention and removal of *Escherichia coli* O157:H7 on surfaces of selected fruits. *Journal of Food Science* 74:E8-E15. 2009.